

References

- [1] I.J. Aalbersberg, G. Rozenberg, and A. Ehrenfeucht. On the membership problem for regular dnlc grammars. *Discrete Appl. Math.*, 13:79–85, 1986.
- [2] A. Iqbal Ali and Jeff L. Kennington. The asymmetric m -travelling salesmen problem: A duality based branch-and-bound algorithm. *Discrete Appl. Math.*, 13:259–276, 1986.
- [3] R.P. Anstee. Invariant sets of arcs in network flow problems. *Discrete Appl. Math.*, 13:1–7, 1986.
- [4] Mikhail J. Atallah and Greg N. Frederickson. A note on finding a maximum empty rectangle. *Discrete Appl. Math.*, 13:87–91, 1986.
- [5] Mikhail J. Atallah and Susanne E. Hambrusch. An assignment algorithm with applications to integrated circuit layout. *Discrete Appl. Math.*, 13:9–22, 1986.
- [6] Francisco Barahona. A solvable case of quadratic 0 – 1-programming. *Discrete Appl. Math.*, 13:23–26, 1986.
- [7] Oscar Barros and Andrés Weintraub. Spatial market equilibrium problems as network models. *Discrete Appl. Math.*, 13:109–130, 1986.
- [8] Jean Bovet. Une amélioration de la méthode de dijkstra pour la recherche d'un plus court chemin dans un réseau. *Discrete Appl. Math.*, 13:93–96, 1986.
- [9] Fred Glover and Beth Novick. The 2-quasi-greedy algorithm for cardinality constrained matroid bases. *Discrete Appl. Math.*, 13:277–286, 1986.
- [10] A.J. Goldman and Prasoon Tiwari. Allowable processing orders in the accelerated cascade algorithm. *Discrete Appl. Math.*, 13:213–221, 1986.
- [11] E. Goles. Antisymmetrical neural networks. *Discrete Appl. Math.*, 13:97–100, 1986.
- [12] Frieda Granot and Rafael Hassin. Multi-terminal maximum flows in node-capacitated networks. *Discrete Appl. Math.*, 13:157–163, 1986.

- [13] S.T. Hedetniemi, Renu Laskar, and John Pfaff. A linear algorithm for finding a minimum dominating set in a cactus. *Discrete Appl. Math.*, 13:287–292, 1986.
- [14] M.S. Hung, W.O. Rom, and A.D. Waren. Degeneracy in transportation problems. *Discrete Appl. Math.*, 13:223–237, 1986.
- [15] Yoshiro Ikura and George L. Nemhauser. Computational experience with a polynomial-time dual simplex algorithm for the transportation problem. *Discrete Appl. Math.*, 13:239–248, 1986.
- [16] Philip C. Jones, Romesh Saigal, and Michael H. Schneider. A variable-dimension homotopy on networks for computing linear spatial equilibria. *Discrete Appl. Math.*, 13:131–156, 1986.
- [17] Darwin D. Klingman and Robert F. Schneider. Microcomputer-based algorithms for large scale shortest path problems. *Discrete Appl. Math.*, 13:183–206, 1986.
- [18] John Mote. The family constrained network problem. *Discrete Appl. Math.*, 13:249–257, 1986.
- [19] Nancy V. Phillips. A new variant of the partitioning shortest path algorithm. *Discrete Appl. Math.*, 13:207–212, 1986.
- [20] Svatopluk Poljak and Daniel Turzík. On an application of convexity to discrete systems. *Discrete Appl. Math.*, 13:27–32, 1986.
- [21] Svatopluk Poljak and Daniel Turzík. On pre-periods of discrete influence systems. *Discrete Appl. Math.*, 13:33–39, 1986.
- [22] I.G. Rosenberg. Structural rigidity ii: Almost infinitesimally rigid bar frameworks. *Discrete Appl. Math.*, 13:41–59, 1986.
- [23] M.T. Shing and T.C. Hu. A decomposition algorithm for multi-terminal network flows. *Discrete Appl. Math.*, 13:165–181, 1986.
- [24] Milan Vlach. Conditions for the existence of solutions of the three-dimensional planar transportation problem. *Discrete Appl. Math.*, 13:61–78, 1986.
- [25] Yechezkel Zalcstein and Stanley P. Franklin. Testing homotopy equivalence is isomorphism complete. *Discrete Appl. Math.*, 13:101–104, 1986.