

References

- [1] Nesrine Abbas and Lorna Stewart. Clustering bipartite and chordal graphs: Complexity, sequential and parallel algorithms. *Discrete Appl. Math.*, 91(1-3):1–23, 1999.
- [2] M. Abellanas, J. García, G. Hernández, M. Noy, and P. Ramos. Bipartite embeddings of trees in the plane. *Discrete Appl. Math.*, 93(2-3):141–148, 1999.
- [3] A.A. Ageev and M.I. Sviridenko. An 0.828-approximation algorithm for the uncapacitated facility location problem. *Discrete Appl. Math.*, 93(2-3):149–156, 1999.
- [4] Rudolf Ahlswede and Zhen Zhang. On maximal shadows of members in left-compressed sets. *Discrete Appl. Math.*, 95(1-3):3–9, 1999.
- [5] Christos Alexopoulos, J. Scott Provan, H. Donald Ratliff, and Bryan R. Stutzman. Minimal connected enclosures on an embedded planar graph. *Discrete Appl. Math.*, 91(1-3):25–38, 1999.
- [6] Noga Alon and Shmuel Onn. Separable partitions. *Discrete Appl. Math.*, 91(1-3):39–51, 1999.
- [7] Charles J. Alpert, Andrew B. Kahng, and So-Zen Yao. Spectral partitioning with multiple eigenvectors. *Discrete Appl. Math.*, 90(1-3):3–26, 1999.
- [8] Martin Anthony. Accuracy of techniques for the logical analysis of data. *Discrete Appl. Math.*, 96-97:247–257, 1999.
- [9] Claudio Arbib, Dario Pacciarelli, and Stefano Smriglio. A three-dimensional matching model for perishable production scheduling. *Discrete Appl. Math.*, 92(1):1–15, 1999.
- [10] T.S. Arthanari and M. Usha. An alternate formulation of the symmetric traveling salesman problem and its properties. *Discrete Appl. Math.*, 98(3):173–190, 1999.
- [11] Luitpold Babel, Andreas Brandstädt, and Van Bang Le. Recognizing the p_4 structure of bipartite graphs. *Discrete Appl. Math.*, 93(2-3):157–168, 1999.

- [12] Luitpold Babel and Stephan Olariu. On the p -connectedness of graphs — a survey. *Discrete Appl. Math.*, 95(1-3):11–33, 1999.
- [13] Jørgen Bang-Jensen, Yubao Guo, and Anders Yeo. A new sufficient condition for a digraph to be hamiltonian. *Discrete Appl. Math.*, 95(1-3):61–72, 1999.
- [14] Jørgen Bang-Jensen and Gregory Gutin. On the complexity of hamiltonian path and cycle problems in certain classes of digraphs. *Discrete Appl. Math.*, 95(1-3):41–60, 1999.
- [15] G. Bär. On the complexity of linear programming in the bss-model. *Discrete Appl. Math.*, 95(1-3):35–40, 1999.
- [16] Vincent Barré and Jean-Luc Fouquet. On minimal imperfect graphs without induced p_5 . *Discrete Appl. Math.*, 94(1-3):9–33, 1999.
- [17] Roberto Battiti and Marco Protasi. Reactive local search techniques for the maximum k -conjunctive constraint satisfaction problem (max- k -ccsp). *Discrete Appl. Math.*, 96 – 97 : 3 – –27, 1999.
- [18] F.E. Bennett. A brief survey of perfect mendelsohn packing covering designs. *Discrete Appl. Math.*, 95(1-3):73–81, 1999.
- [19] C. Benzaken. From logical gates synthesis to chromatic bicritical clutters. *Discrete Appl. Math.*, 96-97:259–305, 1999.
- [20] Jonathan W. Berry and Mark K. Goldberg. Path optimization for graph partitioning problems. *Discrete Appl. Math.*, 90(1-3):27–50, 1999.
- [21] A. Betten and D. Betten. The proper linear spaces on 17 points. *Discrete Appl. Math.*, 95(1-3):83–108, 1999.
- [22] Anton Betten, Reinhard Laue, and Alfred Wassermann. Simple 8 – (40, 11, 1440) designs. *Discrete Appl. Math.*, 95(1-3):109–114, 1999.
- [23] Christian Bey. The erdos-ko-rado bound for the function lattice. *Discrete Appl. Math.*, 95(1-3):115–125, 1999.
- [24] S.L. Bezrukov, R. Elsässer, and U.-P. Schroeder. On k -partitioning of hamming graphs. *Discrete Appl. Math.*, 95(1-3):127–140, 1999.

- [25] Jan C. Bioch, Toshihide Ibaraki, and Kazuhisa Makino. Minimum self-dual decompositions of positive dual-minor boolean functions. *Discrete Appl. Math.*, 96-97:307–326, 1999.
- [26] J. Blazewicz, A. Hertz, D. Kobler, and D. de Werra. On some properties of dna graphs. *Discrete Appl. Math.*, 98(1-2):1–19, 1999.
- [27] Jacek Błażewicz, Maciej Drozdowski, Frédéric Guinand, and Denis Trystram. Scheduling a divisible task in a two-dimensional toroidal mesh. *Discrete Appl. Math.*, 94(1-3):35–50, 1999.
- [28] Zhou Bo and Bolian Liu. Matrices with maximum exponents in the class of doubly stochastic primitive matrices. *Discrete Appl. Math.*, 91(1-3):53–66, 1999.
- [29] Alexander Bockmayr, Friedrich Eisenbrand, Mark Hartmann, and Andreas S. Schulz. On the chvátal rank of polytopes in the 0/1 cube. *Discrete Appl. Math.*, 98(1-2):21–27, 1999.
- [30] Manjit Borah, Robert Michael Owens, and Mary Jane Irwin. A fast and simple steiner routing heuristic. *Discrete Appl. Math.*, 90(1-3):51–67, 1999.
- [31] Richard B. Borie and Jeremy P. Spinrad. Construction of a simple elimination scheme for a chordal comparability graph in linear time. *Discrete Appl. Math.*, 91(1-3):287–292, 1999.
- [32] Endre Boros. Maximum renamable horn sub-cnfs. *Discrete Appl. Math.*, 96-97:29–40, 1999.
- [33] Endre Boros, Peter L. Hammer, Michel Minoux, and Jr. Rader, David J. Optimal cell flipping to minimize channel density in vlsi design and pseudo-boolean optimization. *Discrete Appl. Math.*, 90(1-3):69–88, 1999.
- [34] J.P. Boufflet and J. Carlier. An exact method for minimizing the makespan of an application processed on a master slave bus oriented multiprocessor system. *Discrete Appl. Math.*, 94(1-3):51–76, 1999.
- [35] Ulrik Brandes, Wolfram Schlickenrieder, Gabriele Neyer, Dorothea Wagner, and Karsten Weihe. Planet — a software package of algorithms and heuristics for disjoint paths in *planar networks*. *Discrete Appl. Math.*, 92(2-3):91–110, 1999.

- [36] Andreas Brandstädt and Peter L. Hammer. On the stability number of claw-free p_5 -free and more general graphs. *Discrete Appl. Math.*, 95(1-3):163–167, 1999.
- [37] Andreas Brandtstädt and Van Bang Le. Tree- and forest-perfect graphs. *Discrete Appl. Math.*, 95(1-3):141–162, 1999.
- [38] Peter Brucker, Thomas Hilbig, and Johann Hurink. A branch and bound algorithm for a single-machine scheduling problem with positive and negative time-lags. *Discrete Appl. Math.*, 94(1-3):77–99, 1999.
- [39] E. Rodney Canfield and Konrad Engel. An upper bound for the size of the largest antichain in the poset of partitions of an integer. *Discrete Appl. Math.*, 95(1-3):169–180, 1999.
- [40] Alberto Caprara and Juan José Salazar González. Separating lifted odd-hole inequalities to solve the index selection problem. *Discrete Appl. Math.*, 92(2-3):111–134, 1999.
- [41] Bradley S. Carlson, C.Y. Roger Chen, and Dikran S. Meliksetian. Transistor chaining in static cmos functional cells of arbitrary planar topology. *Discrete Appl. Math.*, 90(1-3):89–114, 1999.
- [42] Soo Y. Chang and Hark-Chin Hwang. The worst-case analysis of the multifit algorithm for scheduling nonsimultaneous parallel machines. *Discrete Appl. Math.*, 92(2-3):135–147, 1999.
- [43] Chandra Chekuri and Rajeev Motwani. Precedence constrained scheduling to minimize sum of weighted completion times on a single machine. *Discrete Appl. Math.*, 98(1-2):29–38, 1999.
- [44] Rachid Chérifi, Sylvain Gravier, Xavier Lagrula, Charles Payan, and Ismaïl Zighed. Domination number of the cross product of paths. *Discrete Appl. Math.*, 94(1-3):101–139, 1999.
- [45] Steve Y. Chiu, Jr. Cox, Louis A., and Xiaorong Sun. Optimal sequential inspections of reliability systems subject to parallel-chain precedence constraints. *Discrete Appl. Math.*, 96-97:327–336, 1999.
- [46] Sunil Chopra and Jonathan H. Owen. A note on formulations for the a -partition problem on hypergraphs. *Discrete Appl. Math.*, 90(1-3):115–133, 1999.

- [47] Philippe Chrétienne. List schedules for cyclic scheduling. *Discrete Appl. Math.*, 94(1-3):141–159, 1999.
- [48] Serafino Cicerone and Gabriele di Stefano. Graph classes between parity and distance-hereditary graphs. *Discrete Appl. Math.*, 95(1-3):197–216, 1999.
- [49] Serafino Cicerone and Gabriele di Stefano. On the extension of bipartite to parity graphs. *Discrete Appl. Math.*, 95(1-3):181–195, 1999.
- [50] Dietmar Cieslik. The steiner ratio of \uparrow_{2k}^d . *Discrete Appl. Math.*, 95(1-3):217–221, 1999.
- [51] Nadia Creignou and Hervé Daude. Satisfiability threshold for random xor-cnf formulas. *Discrete Appl. Math.*, 96-97:41–53, 1999.
- [52] Jürgen Dassow and Gheorghe Păun. On the regularity of languages generated by context-free evolutionary grammars. *Discrete Appl. Math.*, 92(2-3):205–209, 1999.
- [53] Celina M.H. de Figueiredo, John Gimbel, Célia P. Mello, and Jayme L. Szwarcfiter. Even and odd pairs in comparability and in p_4 -comparability graphs. *Discrete Appl. Math.*, 91(1-3):293–297, 1999.
- [54] Caterina de Simone and János Körner. On the odd cycles of normal graphs. *Discrete Appl. Math.*, 94(1-3):161–169, 1999.
- [55] D. de Werra. On a multiconstrained model for chromatic scheduling. *Discrete Appl. Math.*, 94(1-3):171–180, 1999.
- [56] D. de Werra, Ch. Eisenbeis, S. Lelait, and B. Marmol. On a graph-theoretical model for cyclic register allocation. *Discrete Appl. Math.*, 93(2-3):191–203, 1999.
- [57] P. Dell’Olmo and M. Grazia Speranza. Approximation algorithms for partitioning small items in unequal bins to minimize the total size. *Discrete Appl. Math.*, 94(1-3):181–191, 1999.
- [58] Jitender S. Deogun, Ton Kloks, Dieter Kratsch, and Haiko Müller. On the vertex ranking problem for trapezoid, circular-arc and other graphs. *Discrete Appl. Math.*, 98(1-2):39–63, 1999.

- [59] Dominique Désérable. A family of cayley graphs on the hexavalent grid. *Discrete Appl. Math.*, 93(2-3):169–189, 1999.
- [60] Michael J. Dinneen, Jose A. Ventura, Mark C. Wilson, and Golbon Zakeri. Compound constructions of broadcast networks. *Discrete Appl. Math.*, 93(2-3):205–232, 1999.
- [61] Feodor F. Dragan. Almost diameter of a house-hole-free graph in linear time via lexbfs. *Discrete Appl. Math.*, 95(1-3):223–239, 1999.
- [62] Feodor F. Dragan and Falk Nicolai. Lexbfs-orderings of distance-hereditary graphs with application to the diametral pair problem. *Discrete Appl. Math.*, 98(3):191–207, 1999.
- [63] Matthias Ehrgott, Horst W. Hamacher, and Stefan Nickel. Geometric methods to solve max-ordering location problems. *Discrete Appl. Math.*, 93(1):3–20, 1999.
- [64] Thomas Eiter, Toshihide Ibaraki, and Kazuhisa Makino. Bidual horn functions and extensions. *Discrete Appl. Math.*, 96-97:55–88, 1999.
- [65] Oya Ekin, Peter L. Hammer, and Alexander Kogan. On connected boolean functions. *Discrete Appl. Math.*, 96-97:337–362, 1999.
- [66] Ioannis Z. Emiris and Jan Verschelde. How to count efficiently all affine roots of a polynomial system. *Discrete Appl. Math.*, 93(1):21–32, 1999.
- [67] Hikoe Enomoto, Miki Shimabara Miyauchi, and Katsuhiko Ota. Lower bounds for the number of edge-crossings over the spine in a topological book embedding of a graph. *Discrete Appl. Math.*, 92(2-3):149–155, 1999.
- [68] Hazel Everett, Ferran Hurtado, and Marc Noy. Stabbing information of a simple polygon. *Discrete Appl. Math.*, 91(1-3):67–82, 1999.
- [69] Michael J. Fischer and Michael S. Paterson. Optimal layout of edge-weighted forests. *Discrete Appl. Math.*, 90(1-3):135–159, 1999.
- [70] Paul Fischer, Norbert Klasner, and Ingo Wegener. On the cut-off point for combinatorial group testing. *Discrete Appl. Math.*, 91(1-3):83–92, 1999.
- [71] Fedor V. Fomin. Note on a helicopter search problem on graphs. *Discrete Appl. Math.*, 95(1-3):241–249, 1999.

- [72] John Franco, Judy Goldsmith, John Schlipf, Ewald Speckenmeyer, and R.P. Swaminathan. An algorithm for the class of pure implicational formulas. *Discrete Appl. Math.*, 96-97:89–106, 1999.
- [73] Satoru Fujishige and Satoru Iwata. Minimizing a submodular function arising from a concave function. *Discrete Appl. Math.*, 92(2-3):211–215, 1999.
- [74] Ian Gambini. A method for cutting squares into distinct squares. *Discrete Appl. Math.*, 98(1-2):65–80, 1999.
- [75] Joseph L. Ganley. Computing optimal rectilinear steiner trees: A survey and experimental evaluation. *Discrete Appl. Math.*, 90(1-3):161–171, 1999.
- [76] Krzysztof Giaro, Marek Kubale, and Michał Małafiejski. On the deficiency of bipartite graphs. *Discrete Appl. Math.*, 94(1-3):193–203, 1999.
- [77] Andreas Goerdt. A remark on random 2-sat. *Discrete Appl. Math.*, 96-97:107–110, 1999.
- [78] Mordecai Golin and Assaf Schuster. Optimal point-to-point broadcast algorithms via lopsided trees. *Discrete Appl. Math.*, 93(2-3):233–263, 1999.
- [79] L. Gross and G. Farin. A transfinite form of sibson’s interpolant. *Discrete Appl. Math.*, 93(1):33–50, 1999.
- [80] M. Grüttmüller. On the pbd-closure of sets containing 3. *Discrete Appl. Math.*, 95(1-3):251–272, 1999.
- [81] Jun Gu. The multi-sat algorithm. *Discrete Appl. Math.*, 96-97:111–126, 1999.
- [82] Yubao Guo. Outpaths in semicomplete multipartite graphs. *Discrete Appl. Math.*, 95(1-3):273–277, 1999.
- [83] Venkatesan Guruswami. Maximum cut on line and total graphs. *Discrete Appl. Math.*, 92(2-3):217–221, 1999.
- [84] V. Gurvich and L. Khachiyan. On generating the irredundant conjunctive and disjunctive normal forms of monotone boolean functions. *Discrete Appl. Math.*, 96-97:363–373, 1999.
- [85] Susanne E. Hambrusch and Hyeong-Seok Lim. Minimizing broadcast costs under edge reductions in tree networks. *Discrete Appl. Math.*, 91(1-3):93–117, 1999.

- [86] Pierre Hansen, Brigitte Jaumard, and Christoph Meyer. On lower bounds for numbered complete graphs. *Discrete Appl. Math.*, 94(1-3):205–225, 1999.
- [87] Pierre Hansen, Brigitte Jaumard, and Guy-Blaise Douanya Nguetse. Best second order bounds for two-terminal network reliability with dependent edge failures. *Discrete Appl. Math.*, 96-97:375–393, 1999.
- [88] Heiko Harborth and Meinhard Möller. Weakened ramsey numbers. *Discrete Appl. Math.*, 95(1-3):279–284, 1999.
- [89] L.H. Harper. On an isoperimetric problem for hamming graphs. *Discrete Appl. Math.*, 95(1-3):285–309, 1999.
- [90] Sven Hartmann. Asymptotic results on suborthogonal \vec{G} -decompositions of complete digraphs. *Discrete Appl. Math.*, 95(1-3):311–320, 1999.
- [91] Hovhannes A. Harutyunyan and Arthur L. Liestman. More broadcast graphs. *Discrete Appl. Math.*, 98(1-2):81–102, 1999.
- [92] M.C. Hernando, F. Hurtado, A. Márquez, M. Mora, and M. Noy. Geometric tree graphs of points in convex position. *Discrete Appl. Math.*, 93(1):51–66, 1999.
- [93] Alain Hertz. On perfect switching classes. *Discrete Appl. Math.*, 94(1-3):3–7, 1999.
- [94] Peter Heusch. The complexity of the falsifiability problem for pure implicational formulas. *Discrete Appl. Math.*, 96-97:127–138, 1999.
- [95] A.J.W. Hilton and Jr. Johnson, P.D. The hall number, the hall index, and the total hall number of a graph. *Discrete Appl. Math.*, 94(1-3):227–245, 1999.
- [96] Christoph Hipke, Christian Icking, Rolf Klein, and Elmar Langetepe. How to find a point on a line within a fixed distance. *Discrete Appl. Math.*, 93(1):67–73, 1999.
- [97] Chính T. Hoàng. On the disc-structure of perfect graphs i — the co-paw-structure. *Discrete Appl. Math.*, 94(1-3):247–262, 1999.

- [98] F. Hoffmann, K. Kriegel, and C. Wenk. An applied point pattern matching problem: Comparing 2d patterns of protein spots. *Discrete Appl. Math.*, 93(1):75–88, 1999.
- [99] Juha Honkala. On the equivalence problem of context-free and dt0l languages. *Discrete Appl. Math.*, 98(1-2):147–149, 1999.
- [100] J.N. Hooker and M.A. Osorio. Mixed logical-linear programming. *Discrete Appl. Math.*, 96-97:395–442, 1999.
- [101] Peter Horak, Ljiljana Branković, and Mirka Miller. A combinatorial problem in database security. *Discrete Appl. Math.*, 91(1-3):119–126, 1999.
- [102] Edmund Ihler, Gabriele Reich, and Peter Widmayer. Class steiner trees and vlsi-design. *Discrete Appl. Math.*, 90(1-3):173–194, 1999.
- [103] Wilfried Imrich and Sandi Klavžar. Recognizing graphs of acyclic cubical complexes. *Discrete Appl. Math.*, 95(1-3):321–330, 1999.
- [104] Robert W. Irving and David F. Manlove. The b -chromatic number of a graph. *Discrete Appl. Math.*, 91(1-3):127–141, 1999.
- [105] Jeffrey Jackson, Eli Shamir, and Clara Shwartzman. Learning with queries corrupted by classification noise. *Discrete Appl. Math.*, 92(2-3):157–175, 1999.
- [106] Joanna Józefowska, Marek Mika, Rafał Różycki, Grzegorz Waligóra, and Jan Węglarz. Discrete-continuous scheduling to minimize the makespan for power processing rates of jobs. *Discrete Appl. Math.*, 94(1-3):263–285, 1999.
- [107] Dieter Jungnickel and Alexander Pott. Perfect and almost perfect sequences. *Discrete Appl. Math.*, 95(1-3):331–359, 1999.
- [108] Olivier Karch and Thomas Wahl. Relocalization — theory and practice. *Discrete Appl. Math.*, 93(1):89–108, 1999.
- [109] G.B. Khosrovshahi, H.R. Maimani, and R. Torabi. On trades: An update. *Discrete Appl. Math.*, 95(1-3):361–376, 1999.
- [110] Hans Kleine Büning and Theodor Lettmann. Resolution remains hard under equivalence. *Discrete Appl. Math.*, 96-97:139–148, 1999.

- [111] K.M. Koh and E.G. Tay. On optimal orientations of cartesian products with a bipartite graph. *Discrete Appl. Math.*, 98(1-2):103–120, 1999.
- [112] Guy Kortsarz and David Peleg. Approximating the weight of shallow steiner trees. *Discrete Appl. Math.*, 93(2-3):265–285, 1999.
- [113] Svetlana A. Kravchenko. Minimizing the number of late jobs for the two-machine unit-time job-shop scheduling problem. *Discrete Appl. Math.*, 98(3):209–217, 1999.
- [114] O. Kullmann. On a generalization of extended resolution. *Discrete Appl. Math.*, 96-97:149–176, 1999.
- [115] Uwe Leck and Volker Leck. Orthogonal double covers of complete graphs by trees of small diameter. *Discrete Appl. Math.*, 95(1-3):377–388, 1999.
- [116] Jaeun Lee, Moo Young Sohn, and Hye Kyung Kim. A note on graphs with large girth and small minus domination number. *Discrete Appl. Math.*, 91(1-3):299–303, 1999.
- [117] Ming-Chu Li, Derek G. Corneil, and Eric Mendelsohn. Pancyclicity and np -completeness in planar graphs. *Discrete Appl. Math.*, 98(3):219–225, 1999.
- [118] Gary MacGillivray and Min-Li Yu. Generalized partitions of graphs. *Discrete Appl. Math.*, 91(1-3):143–153, 1999.
- [119] Frédéric Maffray and Myriam Preissmann. Sequential colorings and perfect graphs. *Discrete Appl. Math.*, 94(1-3):287–296, 1999.
- [120] A.R. Mahjoub and C. Nocq. On the linear relaxation of the 2-node connected subgraph polytope. *Discrete Appl. Math.*, 95(1-3):389–416, 1999.
- [121] Kazuhisa Makino and Toshihide Ibaraki. Inner-core and outer-core functions of partially defined boolean functions. *Discrete Appl. Math.*, 96-97:443–460, 1999.
- [122] David F. Manlove. On the algorithmic complexity of twelve covering and independence parameters of graphs. *Discrete Appl. Math.*, 91(1-3):155–175, 1999.
- [123] Jacques Mazoyer and Ivan Rapaport. Inducing an order on cellular automata by a grouping operation. *Discrete Appl. Math.*, 91(1-3):177–196, 1999.

- [124] Colin McDiarmid. Pattern minimisation in cutting stock problems. *Discrete Appl. Math.*, 98(1-2):121–130, 1999.
- [125] Ingrid Mengersen and Jörg Oeckermann. Matching-star ramsey sets. *Discrete Appl. Math.*, 95(1-3):417–424, 1999.
- [126] Donatella Merlini, D.G. Rogers, Renzo Sprugnoli, and M. Cecilia Verri. Underdiagonal lattice paths with unrestricted steps. *Discrete Appl. Math.*, 91(1-3):197–213, 1999.
- [127] Masahiro Miyakawa, Akihiro Nozaki, Grant Pogoyan, and Ivo G. Rosenberg. A map from the lower-half of the n -cube onto the $(n-1)$ -cube which preserves intersecting antichains. *Discrete Appl. Math.*, 92(2-3):223–228, 1999.
- [128] Nicolas Moeri. A technique to find multiple motif occurrences in a biomolecular sequence. *Discrete Appl. Math.*, 94(1-3):297–319, 1999.
- [129] Raffaele Mosca. Stable sets in certain p_6 -free graphs. *Discrete Appl. Math.*, 92(2-3):177–191, 1999.
- [130] Vincent Moulton and Mike Steel. Retractions of finite distance functions onto tree metrics. *Discrete Appl. Math.*, 91(1-3):215–233, 1999.
- [131] Jagannathan Narasimhan, Kazuo Nakajima, and Chong S. Rim. A graph theoretical approach for the yield enhancement of reconfigurable vlsi/wsi arrays. *Discrete Appl. Math.*, 90(1-3):195–221, 1999.
- [132] Meei Pyng Ng, Mike Steel, and Nicholas C. Wormald. The difficulty of constructing a leaf-labelled tree including or avoiding given subtrees. *Discrete Appl. Math.*, 98(3):227–235, 1999.
- [133] Kari J. Nurmela and Patric R.J. Östergård. Covering t -sets with $(t+2)$ -sets. *Discrete Appl. Math.*, 95(1-3):425–437, 1999.
- [134] J. Opatrny and D. Sotteau. Embeddings of complete binary trees into grids and extended grids with total vertex-congestion 1. *Discrete Appl. Math.*, 98(3):237–254, 1999.
- [135] Paul Peart and Wen-jin Woan. A divisibility property for a subgroup of riordan matrices. *Discrete Appl. Math.*, 98(3):255–263, 1999.

- [136] Ján Plesník. Constrained weighted matchings and edge coverings in graphs. *Discrete Appl. Math.*, 92(2-3):229–241, 1999.
- [137] Erich Prisner and Jayme L. Szwarcfiter. Recognizing clique graphs of directed and rooted path graphs. *Discrete Appl. Math.*, 94(1-3):321–328, 1999.
- [138] Xiangtong Qi and Fengsheng Tu. Earliness and tardiness scheduling problems on a batch processor. *Discrete Appl. Math.*, 98(1-2):131–145, 1999.
- [139] Mathieu Raffinot. Asymptotic estimation of the average number of terminal states in dawgs. *Discrete Appl. Math.*, 92(2-3):193–203, 1999.
- [140] G. Richomme and P. Séébold. Characterization of test-sets for overlap-free morphisms. *Discrete Appl. Math.*, 98(1-2):151–157, 1999.
- [141] Giovanni Righini and Marco Trubian. Data-dependent bounds for the general and the asymmetric stacker-crane problems. *Discrete Appl. Math.*, 91(1-3):235–242, 1999.
- [142] Irena Rusu. p_4 -domination in minimal imperfect graphs. *Discrete Appl. Math.*, 94(1-3):329–336, 1999.
- [143] Anita Schöbel. Solving restricted line location problems via a dual interpretation. *Discrete Appl. Math.*, 93(1):109–125, 1999.
- [144] Ulrike Schumacher. Suborthogonal double covers of the complete graph by stars. *Discrete Appl. Math.*, 95(1-3):439–444, 1999.
- [145] András Sebő and Zeev Waksman. Optimal binary trees with order constraints. *Discrete Appl. Math.*, 91(1-3):305–311, 1999.
- [146] C.-J. Shi and J.A. Brzozowski. A characterization of signed hypergraphs and its applications to vlsi via minimization and logic synthesis. *Discrete Appl. Math.*, 90(1-3):223–243, 1999.
- [147] R.A. Sitters. A short proof of a conjecture on the t_r -choice number of even cycles. *Discrete Appl. Math.*, 92(2-3):243–246, 1999.
- [148] E. Sojka. A new algorithm for jordan sorting: Its average-case analysis. *Discrete Appl. Math.*, 93(1):127–140, 1999.

- [149] Jan Stallaert. Valid inequalities and separation for capacitated fixed charge flow problems. *Discrete Appl. Math.*, 98(3):265–274, 1999.
- [150] V.A. Strusevich. A heuristic for the two-machine open-shop scheduling problem with transportation times. *Discrete Appl. Math.*, 93(2-3):287–304, 1999.
- [151] L.G. Tallini and U. Vaccaro. Efficient m -ary balanced codes. *Discrete Appl. Math.*, 92(1):17–56, 1999.
- [152] Xuehou Tan and Xiaoyu Song. Routing multiterminal nets on a hexagonal grid. *Discrete Appl. Math.*, 90(1-3):245–255, 1999.
- [153] Feng Tian, Bing Wei, and Lei Zhang. Hamiltonicity in 3-domination-critical graphs with $\alpha = \delta + 2$. *Discrete Appl. Math.*, 92(1):57–70, 1999.
- [154] Peter Tittmann. Partitions and network reliability. *Discrete Appl. Math.*, 95(1-3):445–453, 1999.
- [155] Ioan Tomescu. Some extremal properties of the degree distance of a graph. *Discrete Appl. Math.*, 98(1-2):159–163, 1999.
- [156] Kenji Uemura, Takeo Yaku, and Kimio Sugita. Induced permutation automata and coverings of strongly connected automata. *Discrete Appl. Math.*, 91(1-3):243–249, 1999.
- [157] Alasdair Urquhart. The symmetry rule in propositional logic. *Discrete Appl. Math.*, 96-97:177–193, 1999.
- [158] Allen van Gelder. Complexity analysis of propositional resolution with autarky pruning. *Discrete Appl. Math.*, 96-97:195–221, 1999.
- [159] Stan van Hoesel and Albert Wagelmans. On the complexity of postoptimality analysis of 0/1 programs. *Discrete Appl. Math.*, 91(1-3):251–263, 1999.
- [160] Hans van Maaren. Elliptic approximations of propositional formulae. *Discrete Appl. Math.*, 96-97:223–244, 1999.
- [161] H. Verrall. Compatible hamilton decompositions of directed wrapped butterfly graphs. *Discrete Appl. Math.*, 91(1-3):265–275, 1999.
- [162] Annegret Wagler. Critical edges in perfect line graphs and some polyhedral consequences. *Discrete Appl. Math.*, 95(1-3):455–466, 1999.

- [163] Gerhard J. Woeginger. Sensitivity analysis for knapsack problems: Another negative result. *Discrete Appl. Math.*, 92(2-3):247–251, 1999.
- [164] Henry Wolkowicz and Qing Zhao. Semidefinite programming relaxations for the graph partitioning problem. *Discrete Appl. Math.*, 96-97:461–479, 1999.
- [165] Fuji Zhang and Chang An. Acyclic molecules with greatest homo-lumo separation. *Discrete Appl. Math.*, 98(1-2):165–171, 1999.
- [166] Fuji Zhang and Huaien Li. On acyclic conjugated molecules with minimal energies. *Discrete Appl. Math.*, 92(1):71–84, 1999.
- [167] Shengyuan Zhang. On the nonexistence of extremal self-dual codes. *Discrete Appl. Math.*, 91(1-3):277–286, 1999.