

## References

- [1] J. Apel and W. Lassner. An extension of buchberger's algorithm and calculations in enveloping fields of lie algebras. *J. Symbolic Computation*, 6(2-3):361–370, 1988.
- [2] Leo Bachmair and Nachum Dershowitz. Critical pair criteria for completion. *J. Symbolic Computation*, 6(1):1–18, 1988.
- [3] David Bayer and Ian Morrison. Standard bases and geometric invariant theory — i. initial ideals and state polytopes. *J. Symbolic Computation*, 6(2-3):209–217, 1988.
- [4] David Bayer and Michael Stillman. On the complexity of computing syzygies. *J. Symbolic Computation*, 6(2-3):135–147, 1988.
- [5] Michela Brundu and Fabio Rossi. On the computation of generalized standard bases. *J. Symbolic Computation*, 6(2-3):323–343, 1988.
- [6] Domenico Cantone, Susanna Ghelfo, and Eugenio Omodeo. The automation of syllogistic — i. syllogistic normal forms. *J. Symbolic Computation*, 6(1):83–98, 1988.
- [7] Giuseppa Carrà Ferro. Gröbner bases and hilbert schemes — i. *J. Symbolic Computation*, 6(2-3):219–230, 1988.
- [8] J.H. Davenport. Computer algebra applied to itself. *J. Symbolic Computation*, 6(1):127–132, 1988.
- [9] J. Dixmier and D. Lazard. Minimum number of fundamental invariants for the binary form of degree 7. *J. Symbolic Computation*, 6(1):113–115, 1988.
- [10] Tatiana Gateva-Ivanova and Victor Latyshev. On recognisable properties of associative algebras. *J. Symbolic Computation*, 6(2-3):371–388, 1988.
- [11] Rüdiger Gebauer and H. Michael Möller. On an installation of buchberger's algorithm. *J. Symbolic Computation*, 6(2-3):275–286, 1988.
- [12] Marc Giusti. Combinatorial dimension theory of algebraic varieties. *J. Symbolic Computation*, 6(2-3):249–265, 1988.

- [13] Abdelilah Kandri-Rody and Deepak Kapur. Computing a gröbner basis of a polynomial ideal over a euclidean domain. *J. Symbolic Computation*, 6(1):37–57, 1988.
- [14] Deepak Kapur, David R. Musser, and Paliath Narendran. Only prime superpositions need be considered in the knuth-bendix completion procedure. *J. Symbolic Computation*, 6(1):19–36, 1988.
- [15] Heinz Kredel and Volker Weispfenning. Computing dimension and independent sets for polynomial ideals. *J. Symbolic Computation*, 6(2-3):231–247, 1988.
- [16] Gerhard O. Michler. An algorithm for determining the simplicity of a modular group representation. *J. Symbolic Computation*, 6(1):105–111, 1988.
- [17] A. Miola and T. Mora. Constructive lifting in graded structures: A unified view of buchberger and hensel methods. *J. Symbolic Computation*, 6(2-3):305–322, 1988.
- [18] H. Michael Möller. On the construction of gröbner bases using syzygies. *J. Symbolic Computation*, 6(2-3):345–359, 1988.
- [19] Thomas J. Myers and Maya B. Gokhale. Parallel scheduling of recursively defined arrays. *J. Symbolic Computation*, 6(1):65–81, 1988.
- [20] Alexander Ollongren. On a particular restricted five-body problem — an analysis with computer algebra. *J. Symbolic Computation*, 6(1):117–126, 1988.
- [21] Edmund F. Robertson. Tietze transformations with weighted substring search. *J. Symbolic Computation*, 6(1):59–64, 1988.
- [22] R. Shtokhamer. Lifting canonical algorithms for a ring  $r$  to the ring  $r[x]$ . *J. Symbolic Computation*, 6(2-3):169–181, 1988.
- [23] C. Wrathall. The word problem for free partially commutative groups. *J. Symbolic Computation*, 6(1):99–104, 1988.