

References

- [1] Juan Gerardo Alcazar, Josef Schicho, and Juan Rafael Sendra. A delineability-based method for computing critical sets of algebraic surfaces. *J. Symbolic Computation*, 42(6):678–691, 2007.
- [2] Juan Gerardo Alcazar and Juan Rafael Sendra. Local shape of offsets to algebraic curves. *J. Symbolic Computation*, 42(3):338–351, 2007.
- [3] Michael P. Barnett. Mathscape and molecular integrals. *J. Symbolic Computation*, 42(3):265–289, 2007.
- [4] Leonid Bedratyuk. On complete system of invariants for the binary form of degree 7. *J. Symbolic Computation*, 42(10):935–947, 2007.
- [5] Carlos Beltrán and Luis Miguel Pardo. On the probability distribution of singular varieties of given corank. *J. Symbolic Computation*, 42(1-2):4–29, 2007.
- [6] Isabel Bermejo, Ignacio García-Marco, and Juan José Salazar-González. An algorithm for checking whether the toric ideal of an affine monomial curve is a complete intersection. *J. Symbolic Computation*, 42(10):971–991, 2007.
- [7] Ivo Bleylevens, Ralf Peeters, and Bernard Hanzon. Efficiency improvement in an nd systems approach to polynomial optimization. *J. Symbolic Computation*, 42(1-2):30–53, 2007.
- [8] T. Bogart, A.N. Jensen, D. Speyer, B. Sturmfels, and R.R. Thomas. Computing tropical varieties. *J. Symbolic Computation*, 42(1-2):54–73, 2007.
- [9] Alain Bretto, Alain Faisant, and Luc Gillibert. g -graphs: A new representation of groups. *J. Symbolic Computation*, 42(5):549–560, 2007.
- [10] Manuel Bronstein. Structure theorems for parallel integration. *J. Symbolic Computation*, 42(7):757–769, 2007.
- [11] Massimo Caboara, Sara Faridi, and Peter Selinger. Simplicial cycles and the computation of simplicial trees. *J. Symbolic Computation*, 42(1-2):74–88, 2007.

- [12] Peter Cameron and Natalia Iyudu. Graphs of relations and hilbert series. *J. Symbolic Computation*, 42(11-12):1066–1078, 2007.
- [13] Antonio Campillo, Gert-Martin Greuel, and Christoph Lossen. Equisingular calculations for plane curve singularities. *J. Symbolic Computation*, 42(1-2):89–114, 2007.
- [14] José Manuel Casas, Manuel A. Insua, and Manuel Ladra. Poincaré-birkhoff-witt theorem for leibniz n -algebras. *J. Symbolic Computation*, 42(11-12):1052–1065, 2007.
- [15] Howard Cheng and George Labahn. Output-sensitive modular algorithms for polynomial matrix normal forms. *J. Symbolic Computation*, 42(7):733–750, 2007.
- [16] Conrad Creel and Sam Nelson. Symbolic computation with finite biquandles. *J. Symbolic Computation*, 42(10):992–1000, 2007.
- [17] Raymond Curran and Eduardo Cattani. Restriction of a -discriminants and dual defect toric varieties. *J. Symbolic Computation*, 42(1-2):115–135, 2007.
- [18] Carlos D’Andrea, Hoon Hong, Teresa Krick, and Agnes Szanto. An elementary proof of sylvester’s double sums for subresultants. *J. Symbolic Computation*, 42(3):290–297, 2007.
- [19] Jennifer R. Daniel and Aloysius G. Helminck. Computing the fine structure of real reductive symmetric spaces. *J. Symbolic Computation*, 42(5):497–510, 2007.
- [20] Andreas Dolzmann and Volker Weispfenning. Multiple object semilinear motion planning. *J. Symbolic Computation*, 42(3):324–337, 2007.
- [21] Ioana Dumitriu, Alan Edelman, and Gene Shuman. Mops: Multivariate orthogonal polynomials (symbolically). *J. Symbolic Computation*, 42(6):587–620, 2007.
- [22] Marcin Dumnicki and Witold Jarnicki. New effective bounds on the dimension of a linear system in p^2 . *J. Symbolic Computation*, 42(6):621–635, 2007.

- [23] Eivind Eriksen and Trond Stølen Gustavsen. Computing obstructions for existence of connections on modules. *J. Symbolic Computation*, 42(3):313–323, 2007.
- [24] Gareth A. Evans and Christopher D. Wensley. Complete involutive rewriting systems. *J. Symbolic Computation*, 42(11-12):1034–1051, 2007.
- [25] R.M. Falcón and J. Martín-Morales. Gröbner bases and the number of latin squares related to autotopisms of order ≤ 7 . *J. Symbolic Computation*, 42(11-12):1142–1154, 2007.
- [26] K. Fukuda, A.N. Jensen, N. Lauritzen, and R. Thomas. The generic gröbner walk. *J. Symbolic Computation*, 42(3):298–312, 2007.
- [27] A. Galligo and M. Stillman. On the geometry of parametrized bicubic surfaces. *J. Symbolic Computation*, 42(1-2):136–158, 2007.
- [28] Tatiana Gateva-Ivanova and Shahn Majid. Set-theoretic solutions of the yang-baxter equation, graphs and computations. *J. Symbolic Computation*, 42(11-12):1079–1112, 2007.
- [29] Giulio Genovese. Improving the algorithms of berlekamp and niederreiter for factoring polynomials over finite fields. *J. Symbolic Computation*, 42(1-2):159–177, 2007.
- [30] Edward L. Green and Øyvind Solberg. An algorithmic approach to resolutions. *J. Symbolic Computation*, 42(11-12):1012–1033, 2007.
- [31] A. Hefez and M.E. Hernandes. Standard bases for local rings of branches and their modules of differentials. *J. Symbolic Computation*, 42(1-2):178–191, 2007.
- [32] Xavier Hernández, Josep M. Miret, and Sebastià Xambó-Descamps. Computing the characteristic numbers of the variety of nodal plane cubics in P^3 . *J. Symbolic Computation*, 42(1-2):192–202, 2007. see Erratum in J. Symbolic Computation, Vol. 44, 2009, No. 4, 417-418.
- [33] Hoon Hong and John Perry. Are buchberger’s criteria necessary for the chain condition? *J. Symbolic Computation*, 42(7):717–732, 2007.

- [34] Evelyne Hubert and Irina A. Kogan. Rational invariants of a group action. construction and rewriting. *J. Symbolic Computation*, 42(1-2):203–217, 2007.
- [35] Gabriela Jeronimo and Juan Sabia. Computing multihomogeneous resultants using straight-line programs. *J. Symbolic Computation*, 42(1-2):218–235, 2007.
- [36] Manuel Kauers. Summation algorithms for stirling number identities. *J. Symbolic Computation*, 42(10):948–970, 2007.
- [37] Hitoshi Koshita. An example of relations on the ext-quiver for the suzuki group $sz(8)$ in characteristic 2. *J. Symbolic Computation*, 42(4):429–442, 2007.
- [38] Temur Kutsia. Solving equations with sequence variables and sequence functions. *J. Symbolic Computation*, 42(3):352–388, 2007.
- [39] Chi-Ming Lam, Vladimir Shpilrain, and Jie-Tai Yu. Recognizing and parametrizing curves isomorphic to a line. *J. Symbolic Computation*, 42(7):751–756, 2007.
- [40] Daniel Lazard and Fabrice Rouillier. Solving parametric polynomial systems. *J. Symbolic Computation*, 42(6):636–667, 2007.
- [41] Grégoire Lecerf. Improved dense multivariate polynomial factorization algorithms. *J. Symbolic Computation*, 42(4):477–494, 2007.
- [42] Alexander B. Levin. Gröbner bases with respect to several orderings and multivariable dimension polynomials. *J. Symbolic Computation*, 42(5):561–578, 2007.
- [43] Marc Moreno Maza, Greg Reid, Robin Scott, and Wenyuan Wu. On approximate triangular decompositions in dimension zero. *J. Symbolic Computation*, 42(7):693–716, 2007.
- [44] Gábor P. Nagy and Petr Vojtěchovský. The moufang loops of order 64 and 81. *J. Symbolic Computation*, 42(9):871–883, 2007.
- [45] Franz Pauer. Gröbner bases with coefficients in rings. *J. Symbolic Computation*, 42(11-12):1003–1011, 2007.

- [46] Sonia Pérez-Díaz. Computation of the singularities of parametric plane curves. *J. Symbolic Computation*, 42(8):835–857, 2007.
- [47] Jana Pílníková. Trivializing a central simple algebra of degree 4 over the rational numbers. *J. Symbolic Computation*, 42(6):579–586, 2007.
- [48] Alban Quadrat and Daniel Robertz. Computation of bases of free modules over the weyl algebras. *J. Symbolic Computation*, 42(11-12):1113–1141, 2007.
- [49] E. Rodríguez-Carbonell and D. Kapur. Generating all polynomial invariants in simple loops. *J. Symbolic Computation*, 42(4):443–476, 2007.
- [50] C.J. Sangwin and P. Ramsden. Linear syntax for communicating elementary mathematics. *J. Symbolic Computation*, 42(9):920–934, 2007.
- [51] Philippe Saux Picart and Cyril Brunie. Symmetric subresultants and applications. *J. Symbolic Computation*, 42(9):884–919, 2007.
- [52] Peter Schauenburg. A gröbner-based treatment of elimination theory for affine varieties. *J. Symbolic Computation*, 42(9):859–870, 2007.
- [53] Michael C. Slattery. Generation of groups of square-free order. *J. Symbolic Computation*, 42(6):668–677, 2007.
- [54] Quoc-Nam Tran. A new class of term orders for elimination. *J. Symbolic Computation*, 42(5):533–548, 2007.
- [55] Joris van der Hoeven. Around the numeric-symbolic computation of differential galois groups. *J. Symbolic Computation*, 42(1-2):236–264, 2007.
- [56] Joris van der Hoeven. Efficient accelero-summation of holonomic functions. *J. Symbolic Computation*, 42(4):389–428, 2007.
- [57] Joris van der Hoeven. Generalized power series solutions to linear partial differential equations. *J. Symbolic Computation*, 42(8):771–791, 2007.
- [58] Joris van der Hoeven. New algorithms for relaxed multiplication. *J. Symbolic Computation*, 42(8):792–802, 2007.

- [59] Helena Verrill and David Joyner. Computing with toric varieties. *J. Symbolic Computation*, 42(5):511–532, 2007.
- [60] Michael Wibmer. Gröbner bases for families of affine or projective schemes. *J. Symbolic Computation*, 42(8):803–834, 2007.